

MEDICAL INFORMATION

4. Zimmerman HJ: Aspirin-induced hepatic injury. *Ann Intern Med* 80:103-105, Jan 1974
5. Seaman WE, Plotz PH: Effect of aspirin on liver tests in patients with RA or SLE and in normal volunteers. *Arthritis Rheum* 19:155-160, Mar 1976
6. Saltzman DA, Gall EP, Robinson SF: Aspirin-induced hepatic dysfunction in a patient with adult rheumatoid arthritis. *Am J Dig Dis* 21:815-820, Sep 1976
7. Miller JJ, Weissman DB: Correlations between transaminase concentrations and serum salicylate concentration in juvenile rheumatoid arthritis. *Arthritis Rheum* 19:115-118, Jan 1976
8. O'Gorman T, Koff RS: Salicylate hepatitis. *Gastroenterology* 72:726-728, Apr 1977
9. Barker JD, deCarle DJ, Anuras S: Chronic excessive acetaminophen use and liver damage. *Ann Intern Med* 87:299-301, Sep 1977
10. Johnson CK, Tolman KG: Chronic liver disease and acetaminophen. *Ann Intern Med* 87:302-304, Sep 1977
11. Mitchell JR: Host susceptibility and acetaminophen liver injury. *Ann Intern Med* 87:377-378, Sep 1977
12. Drivsholm AA, Madsen S: The influence of treatment with sodium salicylate on the serum glutamic oxalacetic transaminase activity. *Scand J Clin Lab Invest* 13:442-446, 1961
13. Woodbury DM, Fingl E: Analgesic-antipyretics, anti-inflammatory agents, and drugs employed in the therapy of gout, Chap 17, *In Goodman CS, Gilman A (Eds): The Pharmacological Basis of Therapeutics*, 5th Ed. New York, Macmillan, 1975
14. Manso C, Taranta A, Nydick I: Effect of aspirin administration on serum glutamic oxalacetic and glutamic pyruvic transaminases in children. *Proc Soc Exp Biol Med* 93:84-88, Oct 1956
15. Iancu T: Serum transaminases and salicylate therapy. *Br Med J* 2:167, Apr 1972
16. Ameer B, Greenblatt DJ: Acetaminophen. *Ann Intern Med* 87:202-209, Aug 1977
17. Editorial: Paracetamol hepatotoxicity. *Lancet* 2:1189-1191, Dec 1975
18. Ferguson DR, Snyder SK, Cameron AJ: Hepatotoxicity in acetaminophen poisoning. *Mayo Clin Proc* 52:246-248, Apr 1977
19. Rosenberg DM, Neelon FA: Acetaminophen and liver disease. *Ann Intern Med* 88:129, Jan 1978
20. Mitchell JR, Thorgenson SS, Potter WZ, et al: Acetaminophen-induced hepatic injury: Protective role of glutathione in man and rationale for therapy. *Clin Pharmacol Therap* 16:676-684, Oct 1974
21. Wright N, Prescott LF: Potentiation by previous drug therapy of hepatotoxicity following paracetamol overdosage. *Scott Med J* 18:56-58, Jan 1973

Refer to: Rosenblatt S, Hershman JM: Serum thyroid hormones in patients on replacement or suppressive doses of L-thyroxine. (Medical Information). *West J Med* 129:165-167, Aug 1978

Medical Information

Serum Thyroid Hormones in Patients on Replacement or Suppressive Doses of L-Thyroxine

SIDNEY ROSENBLATT, MD
JEROME M. HERSHMAN, MD
Los Angeles

CHANGES IN THE normal circulating levels of thyroid hormones occur in patients taking replacement or suppressive doses of L-thyroxine (T_4). Most of the circulating pool of triiodothyronine (T_3) is derived from peripheral monodeiodination of thyroxine.¹⁻³ Adequate treatment with L-thyroxine usually produces serum T_4 levels similar to or slightly higher than control values, T_3 levels

From the Endocrinology Section, VA Wadsworth Medical Center, Los Angeles, and the Department of Medicine, University of California, Los Angeles, School of Medicine.

Supported by the Medical Research Service of the Veterans Administration.

Reprint requests to Sidney Rosenblatt, MD, Endocrinology Section, Wadsworth VA Hospital, Wilshire and Sawtelle Blvds., Los Angeles, CA 90073.

ABBREVIATIONS USED IN TEXT

T_3 = triiodothyronine
 T_4 = L-thyroxine
TSH = thyroid-stimulating hormone

are often slightly lower than controls, and the T_4/T_3 ratio would be expected to be higher than control values. We measured serum concentrations of T_4 , T_3 and thyroid-stimulating hormone (TSH) in patients with a variety of thyroid disorders to assess the significance of changes in circulating thyroid hormone levels in patients taking L-thyroxine.

Methods

A retrospective analysis was made of 132 patients receiving adequate suppressive or replacement doses of L-thyroxine. A total of 71 patients were in the hypothyroid group, and 61 were in the euthyroid group being treated for nodular or diffuse goiter. Patients with autonomously functioning nodular goiters were not included. The dose range was 0.1 to 0.3 mg of L-thyroxine, and adequate treatment was assessed by normal concentrations of serum thyrotropin (less than 5 μ U/per ml) in all patients without pituitary disease. The patients with pituitary disease all had normal levels of serum T_3 and T_4 . Serum T_3 , T_4 and TSH were measured by radioimmunoassays.⁴ All patients were receiving therapy for at least three months. The patients were otherwise

healthy which excluded non-thyroidal alterations in serum thyroid hormones.

A total of 161 patients without thyroid disease served as normal controls. The age patterns in the controls and treated groups were similar. Statistical significance was determined by Student's t-test.

Results

Table 1 shows the serum T_3 , T_4 , T_4/T_3 ratio and TSH in the control group and the groups treated with various doses of L-thyroxine. Serum T_4 concentrations and the T_4/T_3 ratio were higher in all the treatment groups irrespective of dose ($p < 0.05$). Serum T_3 concentrations were elevated significantly only in the euthyroid treated patients. Serum TSH was undetectable (less than $0.3 \mu\text{U}/\text{per ml}$) in 58 percent of all patients treated for primary hypothyroidism and 72 percent of all patients in the euthyroid treated group.

When patients in both treatment groups were selected because their serum T_4 was in the lower half of the normal range (5 to $8.5 \mu\text{g per dl}$), the T_4/T_3 ratio was 51.4 in the hypothyroid and 48.0 in the euthyroid treated groups. These values were similar to the ratio in untreated controls. The mean T_3 level was $143 \pm 29 \text{ ng per dl}$ in the hypothyroid treated groups and $150 \pm 26 \text{ ng per dl}$ in the euthyroid treated groups, which are similar to T_3 concentrations in controls.

Discussion

The data show that T_4 concentrations and the T_4/T_3 ratio in patients receiving replacement or suppressive doses of L-thyroxine are increased. T_3 levels are normal or increased. Higher levels of thyroid hormones were observed in patients treated for goiter suppression because the therapeutic goal was suppression of endogenous TSH secretion.

In patients taking the usually prescribed doses of L-thyroxine, the T_4/T_3 ratio may be altered for two reasons. Firstly, the serum T_4 level is elevated, suggesting that these doses of L-thyroxine are too high. The mean T_4 level in the hypothyroid group was $2.6 \mu\text{g per dl}$ higher than the mean control value of $7.6 \mu\text{g per dl}$, and the mean T_4 value in the euthyroid group was $3.7 \mu\text{g per dl}$ higher than the controls. To further elucidate this point, we selected those patients receiving therapy who had T_4 levels between 5 and $8.5 \mu\text{g per dl}$. Their T_3 levels and T_4/T_3 ratios were unchanged from those of controls. Therefore, the alterations in thyroid hormone relationships is primarily affected by the serum T_4 level. Second, the circulating levels of T_3 in patients taking L-thyroxine is derived primarily from peripheral monodeiodination.¹⁻³ The reduced thyroidal contribution of T_3 , secondary to suppression or loss of thyroid secretion, probably also plays a role.

TABLE 1.—Serum Thyroid Hormone Concentrations (Mean \pm S.D.)

GROUP	T_4 dose mg	No.	T_4 $\mu\text{g}/\text{dl}$	T_3 ng/dl	T_4/T_3	TSH $\mu\text{U}/\text{ml}$	Undetectable TSH (Percent)
Normal	0	161	7.6 ± 1.6	153.7 ± 32.7	50.8 ± 13.1	1.8 ± 1.6	4.3
Hypothyroid	0.1-0.15	34	9.2* ± 2.4	168.8 ± 64.4	59.4* ± 23.2	1.3* ± 1.6	44†
Hypothyroid	0.2	27	11.3† ± 4.6	173.6 ± 44.0	64.3* ± 17.1	1.1* ± 2.3	66†
Hypothyroid	0.25-0.30	10	10.6† ± 3.4	175.1 ± 22.2	59.3* ± 12.3	0.3† ± 0.8	89†
Euthyroid	0.1-0.15	11	10.0† ± 3.2	173.7* ± 64.8	59.7* ± 13.2	0.5† ± 0.8	63
Euthyroid	0.20	34	11.1† ± 3.3	181.5* ± 45.5	63.2* ± 15.0	0.4† ± 0.7	71
Euthyroid	0.25-0.30	16	12.7† ± 3.0	186.3* ± 49.0	70.1* ± 14.5	0.1† ± 0.2	81
All hypothyroid		71	10.2† ± 3.5	171.5 ± 52.4	61.2* ± 19.7	1.1* ± 2.1	58
All euthyroid		61	11.3† ± 3.3	181.4* ± 49.5	64.4* ± 14.8	0.3† ± 0.6	72

T_3 = triiodothyronine

T_4 = L-thyroxine

TSH = thyroid-stimulating hormone

* $P < 0.05$ compared with normal untreated subjects

† $P < 0.005$

‡Patients with primary hypothyroidism only

MEDICAL INFORMATION

By using a sensitive radioimmunoassay for TSH, we showed that most patients taking L-thyroxine had undetectable levels. Nearly all normal and untreated euthyroid goitrous patients have detectable circulating levels of TSH with this assay. We agree with Stock and co-workers⁵ that the current textbook recommendations for replacement therapy with L-thyroxine are too high. We recommend 0.1 to 0.2 mg of L-thyroxine for replacement therapy. Suppression therapy with this dose range will effectively reduce TSH to undetectable levels in most patients; however, the dose must be individually tailored for each patient.

Summary

Serum thyroid hormones were measured in patients taking L-thyroxine for replacement or suppressive therapy. Their serum T_4 levels and T_4/T_3 ratios were greater than controls and their serum

T_3 levels were normal or slightly increased. This finding was more evident in the euthyroid goitrous group because higher doses of thyroxine were used to suppress TSH secretion. Serum TSH levels were suppressed to undetectable levels in most patients. Replacement therapy for hypothyroidism can be achieved with 0.1 to 0.2 mg of L-thyroxine which will suppress serum TSH to undetectable levels in most patients.

REFERENCES

1. Braverman LE, Ingbar SH, Sterling K: Conversion of thyroxine (T_4) to triiodothyronine (T_3) in athyreotic human subjects. *J Clin Invest* 49:855-864, May 1970
2. Braverman LE, Vagenakis A, Downs P, et al: Effects of replacement doses of sodium-L-thyroxine on the peripheral metabolism of thyroxine to triiodothyronine in man. *J Clin Invest* 52:1010-1017, May 1973
3. Chopra IJ: An assessment of daily production and significance of thyroidal secretion of 3,3',5'-triiodothyronine (reverse T_3) in man. *J Clin Invest* 58:32-40, Jul 1976
4. Azukizawa M, Pekary AE, Hershman JM, et al: Plasma thyrotropin, thyroxine, and triiodothyronine relationships in man. *J Clin Endocrinol Metab* 43:533-542, Sep 1976
5. Stock JM, Surks MI, Oppenheimer JH: Replacement dosage of L-thyroxine in hypothyroidism. *N Engl J Med* 290:529-533, Mar 1974

Herpes Zoster Keratouveitis

YOU SHOULD ALL REMEMBER that those who have herpes zoster keratouveitis in the older age group may very well have an underlying malignancy, and that should be looked for.

—PETER R. LAIBSON, MD, Philadelphia

Extracted from *Audio-Digest Ophthalmology*, Vol. 15, No. 16, in the Audio-Digest Foundation's subscription series of tape-recorded programs. For subscription information: 1577 East Chevy Chase Drive, Glendale, CA 91206.